

# The Role of a Dopamine Stimulated G<sub>q</sub> Pathway in the Development of Striatal Medium Spiny Neurons


Mason Fellmeth, Justin Campagne, Lorene Lanier  
Department of Neuroscience, University of Minnesota – Twin Cities

## Abstract

There exists considerable evidence linking the role of dopamine receptor activation of Medium Spiny Neurons (MSNs) in the striatum to neurite development. A better understanding of this link between receptor activation and dendrite growth and plasticity can significantly contribute towards the understanding of ailments including Huntington's disease, depression, and addiction. **MSNs cultured in the presence of dopamine (+DA) show significant dendrite growth and arborization compared to controls.** Interestingly, cultures treated with dopamine stimulatory (D1) and dopamine inhibitor (D2) specific receptor agonists show little to no change in dendrite development compared to -DA controls. These results implicate an alternative dopamine mechanism separate from that of the traditional D1 and D2 pathways. Previous work has shown that the use of a G<sub>q</sub> coupled Designer Receptor Exclusively Activated by Designer Drugs (DREADD), selectively activated by clozapine-n-oxide (CNO), exhibits dendrite growth similar to that observed in +DA MSNs. In our current study, we tested the potential role of a G<sub>q</sub> coupled dopamine receptor in MSN development using the D1-D2 selective agonist SKF83959.

## Background



- The striatum is a subnucleus of the basal ganglia with the primary cell population consisting of Medium Spiny Neurons (MSNs)<sup>4</sup>
- Abnormal striatal functioning has been linked to ailments including Parkinson's disease, Huntington's disease, and addiction<sup>1,2,8</sup>
- MSNs receive both dopamine and glutamate innervation from the cortex and midbrain respectively<sup>6,7</sup>
- The prevailing model of MSNs shows them as expressing either D1 or D2 dopamine receptors<sup>5</sup>
  - D1 pathway – G<sub>s</sub>; Activates adenylyl cyclase
  - D2 pathway – G<sub>i</sub>; Blocks adenylyl cyclase



## Hypothesis

- If MSN treatment with a dopamine D1 and D2 receptor agonists fails to replicate the dendrite growth observed when cells are cultured with dopamine, then dopamine is likely acting through a different mechanism
  - The activation of a transfected G<sub>q</sub> pathway will mimic results of cells cultured in the presence of dopamine
  - MSNs treated with the D1-D2 receptor agonist SKF83959 will mirror results gained from transfected G<sub>q</sub> treatment and cells cultured with dopamine

## Methods



## Results




Figure 1: Most dendrite arborization occurred between 13-19 div and peaked around ring number 5 and 6 (50-60um from the soma). For 19 div cultures treated with dopamine (+DA), there was a significant increase in ring crossings, signifying increased arborization<sup>3</sup>.




Figure 2: MSNs cultured with D1 receptor agonist (SKF) and/or D2 receptor agonist (quinprole) failed to replicate the increase in arborization observed in cultures treated with dopamine<sup>3</sup>. Suggesting that dopamine is acting via a non-traditional pathway.




Figure 3: The effects of dopamine on dendritic arborization in MSN cell cultures was significantly decreased in the presence of the phospholipase-C antagonist U73122<sup>3</sup>. This suggests the role of a G<sub>q</sub> pathway.





Figure 4: When MSNs were transfected with a designer G<sub>q</sub> receptor specifically activated by CNO (DREADDs), there was a significant increase in dendrite arborization that was comparable to cells that received dopamine treatment. These cells also had significantly greater arborization than cells not exposed to dopamine<sup>9</sup>.



## Acknowledgements and References

- Deuth, A. Y., Colbran, R. J., Winder, D. J., (2007). Striatal plasticity and medium spiny neuron dendritic remodeling in parkinsonism. *Parkinsonism & Related Disorders*, 13, S251-S253
- Glass, M., Dragunow, M., & Faull, R. L. (2000). The pattern of neurodegeneration in Huntington's disease: a comparative study of cannabinoid, dopamine, adenosine and GABA(A) receptor alterations in the human basal ganglia in Huntington's disease. *Neuroscience*, 97(3), 505-19. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/10828533>
- Penrod, R. D., Campagna, J., Panneck, T., Preese, L., & Lanier, L. M. (2015). The presence of cortical neurons in striatal-cortical co-cultures alters the effects of dopamine and BDNF on medium spiny neuron dendritic development. *Frontiers in Cellular Neuroscience*, 9, 269. <https://doi.org/10.3389/fncel.2015.00269>
- Rafols, J. A., Cheng, H. W., & McNeill, T. H. (1989). Golgi study of the mouse striatum: age-related dendritic changes in different neuronal populations. *The Journal of Comparative Neurology*, 2, 2
- Schmidt, U., Pilgrim, C., & Beyer, C. (1998). Differentiative Effects of Dopamine on Striatal Neurons Involve Stimulation of the cAMP/PKA Pathway. *Molecular and Cellular Neuroscience*, 11(1), 9-18. <https://doi.org/10.1006/mcne.1998.0668>
- Swanson, L. W. (1982). The projections of the ventral tegmental area and adjacent regions: A combined fluorescent retrograde tracer and immunofluorescence study in the rat. *Brain Research Bulletin*, 9(1), 321-353. [https://doi.org/10.1016/0361-9230\(82\)90145-9](https://doi.org/10.1016/0361-9230(82)90145-9)
- Wall, N. R., De La Parra, M., Callaway, E. M., & Kretzler, A. C. (2013). Differential Innervation of Direct- and Indirect-Pathway Striatal Projection Neurons. *Neuron*, 79(2), 347-360. <https://doi.org/10.1016/j.neuron.2013.05.01>
- Yager, L. M., Garcia, A. F., Wunsch, A. M., & Ferguson, S. M. (2015). The ins and outs of the striatum: role in drug addiction. *Neuroscience*, 301, 529-41. <https://doi.org/10.1016/j.neurosci>
- UNPUBLISHED – Lanier, L., Campagne, J.